

# The Proportion of SARS-CoV-2 Infections That Are Asymptomatic

## A Systematic Review

Daniel P. Oran, AM, and Eric J. Topol, MD

**Background:** Asymptomatic infection seems to be a notable feature of severe acute respiratory syndrome coronavirus 2 (SARS CoV 2), the pathogen that causes coronavirus disease 2019 (COVID 19), but the prevalence is uncertain.

**Purpose:** To estimate the proportion of persons infected with SARS CoV 2 who never develop symptoms.

**Data Sources:** Searches of Google News, Google Scholar, medRxiv, and PubMed using the keywords *antibodies, asymptomatic, coronavirus, COVID 19, PCR, seroprevalence, and SARS CoV 2*.

**Study Selection:** Observational, descriptive studies and reports of mass screening for SARS CoV 2 that were either cross sectional or longitudinal in design; were published through 17 November 2020; and involved SARS CoV 2 nucleic acid or antibody testing of a target population, regardless of current symptomatic status, over a defined period.

**Data Extraction:** The authors collaboratively extracted data on the study design, type of testing performed, number of participants, criteria for determining symptomatic status, testing results, and setting.

**Data Synthesis:** Sixty one eligible studies and reports were identified, of which 43 used polymerase chain reaction (PCR) testing of nasopharyngeal swabs to detect current SARS

CoV 2 infection and 18 used antibody testing to detect current or prior infection. In the 14 studies with longitudinal data that reported information on the evolution of symptomatic status, nearly three quarters of persons who tested positive but had no symptoms at the time of testing remained asymptomatic. The highest quality evidence comes from nationwide, representative serosurveys of England ( $n = 365\,104$ ) and Spain ( $n = 61\,075$ ), which suggest that at least one third of SARS CoV 2 infections are asymptomatic.

**Limitation:** For PCR-based studies, data are limited to distinguish presymptomatic from asymptomatic infection. Heterogeneity precluded formal quantitative syntheses.

**Conclusion:** Available data suggest that at least one third of SARS CoV 2 infections are asymptomatic. Longitudinal studies suggest that nearly three quarters of persons who receive a positive PCR test result but have no symptoms at the time of testing will remain asymptomatic. Control strategies for COVID 19 should be altered, taking into account the prevalence and transmission risk of asymptomatic SARS CoV 2 infection.

**Primary Funding Source:** National Institutes of Health.

*Ann Intern Med.* 2021;174:655-662. doi:10.7326/M20-6976 **Annals.org**

For author, article, and disclosure information, see end of text.

This article was published at Annals.org on 22 January 2021.

The asymptomatic fraction of infection is the proportion of infected persons who never develop, perceive, and report symptoms (1). Among common pathogens, the asymptomatic fraction varies widely. For example, an asymptomatic carrier state has not been documented for measles virus infection (2), whereas a significant proportion of persons with cytomegalovirus or poliovirus infection have no symptoms and are unaware of infection (3, 4). The asymptomatic fraction of severe acute respiratory syndrome coronavirus 2 (SARS CoV 2) infection seems to be sizable (5). The range of severity of illness associated with SARS CoV 2 infection is noteworthy because it spans asymptomatic infection; mild illness; and severe, life threatening illness.

Perhaps because of this broad spectrum of presentation, the topic of asymptomatic SARS CoV 2 infection has generated some controversy (6). Imprecise use of the term "asymptomatic" is partly to blame. "Asymptomatic" should be reserved for persons who never develop symptoms, whereas "presymptomatic" is a better description of those who have no symptoms when they receive a positive test result but who eventually develop symptoms. We know for certain who is asymptomatic only in retrospect. On the basis of our current knowledge of the natural history of coronavirus disease 2019 (COVID 19), after a person is infected with SARS CoV 2, we must wait approximately 14

days to determine whether symptoms have developed (7). Infection without symptoms, whether presymptomatic or asymptomatic, is important because infected persons can transmit the virus to others even if they have no symptoms (8, 9).

In June 2020, we published a review of the limited data then available on the prevalence of asymptomatic SARS CoV 2 infection (5). Since then, considerable new data have become available. The present review summarizes currently available data that might allow us to estimate the proportion of persons infected with SARS CoV 2 who are asymptomatic.

## METHODS

### Data Sources, Search Terms, and Study Selection

Using the keywords *antibodies, asymptomatic, coronavirus, COVID 19, PCR, seroprevalence, and SARS CoV 2*, we periodically searched Google News, Google Scholar, medRxiv, and PubMed for observational, descriptive studies and reports of mass screening for SARS CoV 2 that were either cross sectional or longitudinal in design; were published through 17 November 2020; and involved SARS CoV 2 nucleic acid or antibody testing of a target population, regardless of current symptomatic status, over a defined period.

### Data Extraction and Quality Assessment

We recorded the total number of persons tested, the number that tested positive, the number of positive cases without symptoms, the criteria for determining symptomatic status, whether the data were cross sectional or longitudinal in nature, whether random selection techniques were used to achieve a representative sample of a target population, and whether the testing involved polymerase chain reaction (PCR) analysis of a nasopharyngeal swab or serologic analysis of antibodies in a blood sample. For longitudinal studies that provided information on the evolution of symptomatic status, we recorded the proportion of persons who tested positive but had no symptoms at the time of testing and who then remained asymptomatic during a follow up period. In addition, we flagged studies that required clarification of ambiguous details.

Studies or reports that are based on PCR results and include only cross sectional data do not make it possible to distinguish between presymptomatic and asymptomatic SARS CoV 2 infection because symptomatic status is observed on only 1 occasion, which may occur before the development of symptoms, if any. In contrast, we can distinguish between presymptomatic and asymptomatic infection with either antibody based studies, in which an interview or questionnaire gathers information about symptoms reported at the time a blood sample is taken and during a prior period, or PCR based studies that include longitudinal data.

In assessing quality, we put the greatest emphasis on random selection of participants to achieve a representative sample of a regional or national population, a large number of study participants ( $n > 10\,000$ ), and study designs that make it possible to distinguish between presymptomatic and asymptomatic infection. Evaluated in this manner, the highest quality evidence comes from large scale, national studies with representative samples that include data from either antibody or longitudinal PCR testing. In **Tables 1** and **2**, we show in boldface the details that increase a study's likelihood of providing higher quality evidence.

### Data Synthesis and Analysis

We synthesized evidence qualitatively by evaluating study design, including whether data were collected longitudinally; testing methods; number of participants; and setting. We compared the range and consistency of estimates of the proportion of persons who tested positive but had no symptoms at the time of testing.

### Role of the Funding Source

The National Institutes of Health played no role in the design, conduct, or analysis of this review or in the decision to submit the manuscript for publication.

## RESULTS

We identified 61 studies or reports that met eligibility criteria. **Table 1** (10–54) summarizes data from the 43 that used PCR testing, and **Table 2** (55–72) summarizes data from the 18 that used antibody testing. The

heterogeneity of the studies in particular, disparate settings and populations precluded quantitative summaries using meta analysis. We summarize the evidence in terms of the number of studies and the range, median, and interquartile range (IQR) for persons who tested positive but had no symptoms at the time of PCR testing or who reported having had no symptoms before or at the time of antibody testing. Thirty of the studies included a list of specific symptoms, independent of signs, used to determine symptomatic status (10–14, 17, 18, 22–28, 35, 36, 38, 42, 49, 51, 55–57, 60–62, 64). Many of the remaining studies used some variation of the catch all phrase “symptoms compatible with COVID 19.”

### Nucleic Acid PCR Testing

Among the 43 studies using PCR testing (10–54), the proportion of persons who tested positive but had no symptoms at the time of testing ranged from 6.3% to 100%, with a median of 65.9% (IQR, 42.8% to 87.0%).

Nineteen of the PCR based studies collected data on symptoms longitudinally after testing, making it possible to distinguish between presymptomatic and asymptomatic infection (15, 17, 18, 20, 22, 25, 26, 27, 32, 37–40, 45, 47, 48, 51, 53, 54). The follow up period in these studies ranged from 2 to 70 days, with a median of 14 days (IQR, 14.0 to 15.8 days). The proportion of persons who tested positive and remained asymptomatic ranged from 6.3% to 91.7%, with a median of 42.5% (IQR, 29.6% to 77.8%).

Of the 19 longitudinal studies, 14 provided information on the evolution of symptomatic status (**Table 3**) (15, 17, 18, 20, 22, 32, 37–40, 47, 51, 53, 54). Among persons who tested positive but had no symptoms at the time of testing, the proportion who remained asymptomatic during a follow up period ranged from 11.1% to 100%, with a median of 72.3% (IQR, 56.7% to 89.7%).

Of the 43 studies that used PCR testing, 24 collected cross sectional data and reported only the symptomatic status at the time of testing, so we could not distinguish between presymptomatic and asymptomatic cases (10–14, 16, 19, 21, 23, 24, 28–31, 33–36, 41–44, 46, 49, 50, 52). In these studies, the proportion of persons who tested positive but had no symptoms at the time of testing ranged from 40.7% to 100%, with a median of 75.5% (IQR, 50.3% to 86.2%).

Of the 43 studies that used PCR testing, 4 used random selection of participants to achieve a representative sample of their target population: residents of England (10–12, 14), Iceland (16), or Indiana (23). Proportions of persons who tested positive but had no symptoms at the time of testing ranged from 43.0% to 76.5%, with a median of 45.6% (IQR, 43.6% to 61.8%). None of the PCR testing studies that used random selection of participants collected longitudinal data on symptoms, so we could not distinguish between presymptomatic and asymptomatic cases.

The largest of the representative data sets, and the largest study identified in our search, was from the REACT (Real time Assessment of Community Transmission) program. REACT has implemented nationwide nucleic acid and antibody testing (discussed later) for SARS CoV 2 of

persons in England aged 5 years and older in multiple phases since May 2020 (10-12). In Table 1, we have combined the results of 6 phases of nucleic acid testing from REACT, yielding data for 932 072 persons (England residents 1). At the time of testing, 1425 of 3029 persons (47.0%) who tested positive had no symptoms. The study did not collect longitudinal data on symptoms, so we could not distinguish between presymptomatic and asymptomatic cases.

The second largest of the representative studies was also from England; it included 36 061 persons tested between 26 April and 27 June 2020 (14). The proportion of persons who tested positive was 0.3%, identical to that reported by REACT, but the proportion of persons who tested positive but had no symptoms at the time of

testing was 74.8%, much larger than in the REACT study. The study did not collect longitudinal data on symptoms, so we could not distinguish between presymptomatic and asymptomatic cases.

In the cross sectional study of Belgian long term care facilities (n = 280 427), age did not seem to affect the proportion of persons who tested positive but had no symptoms at the time of testing (13). The study tested 138 327 staff and 142 100 residents. Median age was 42 years for staff and 85 years for residents; despite this considerable difference, the proportion of those who tested positive without symptoms was 74.0% for staff and 75.3% for residents. This finding is consonant with the finding of a longitudinal study from Vo', Italy, in which more than 85% of the town's 3275 residents were tested:

Table 1. Nucleic Acid PCR Testing

Study or Report	Tested, n*	Longitudinal Data*	Random Sampling*	SARS-CoV-2-Positive, n (%)	Positive, but No Symptoms, n (%)
England residents 1 (10-12)	<b>932 072</b>	No	<b>Yes</b>	3029 (0.3)	1425 (47.0)
Belgium long-term care facility residents and staff (13)	<b>280 427</b>	No	No	8343 (3.0)	6244 (74.8)
England residents 2 (14)	<b>36 061</b>	No	<b>Yes</b>	115 (0.3)	88 (76.5)
U.S. skilled-nursing facility residents (15)†	<b>22 368</b>	<b>Yes</b>	No	5403 (24.2)	2194 (40.6)
Iceland residents (16)	<b>13 080</b>	No	<b>Yes</b>	100 (0.8)	43 (43.0)
Vo', Italy, residents (17)	5155	<b>Yes</b>	No	102 (2.0)	34 (42.5)
U.S. Navy aircraft carrier crew (18)	4779	<b>Yes</b>	No	1271 (26.6)	572 (45.0)
Arkansas, North Carolina, Ohio, and Virginia inmates (19)	4693	No	No	3277 (69.8)	3146 (96.0)
San Francisco, California, residents (20)	3871	<b>Yes</b>	No	83 (2.1)	23 (27.7)
Arkansas poultry plant employees (21)	3748	No	No	481 (12.8)	455 (94.6)
Diamond Princess cruise ship passengers and crew (22)	3618	<b>Yes</b>	No	712 (19.7)	311 (43.7)
Indiana residents (23)‡	3605	No	<b>Yes</b>	47 (1.7)	18 (44.2)
South London, England, nursing home residents and staff (24)	2455	No	No	160 (6.5)	115 (71.9)
U.S. Marine recruits (25)	1801	<b>Yes</b>	No	51 (2.8)	46 (90.2)
Charles de Gaulle aircraft carrier crew (26)	1568	<b>Yes</b>	No	1001 (63.8)	130 (13.0)
Marseille, France, long-term care facility residents (27)	1691	<b>Yes</b>	No	226 (13.4)	46 (23.0)
King County, Washington, homeless shelter residents and staff (28)	1434	No	No	29 (2.0)	21 (72.4)
Germany oncology clinic patients (29)	1286	No	No	40 (3.1)	37 (92.5)
Pasadena, California, long-term care facilities residents and staff (30)	938	No	No	631 (67.3)	257 (40.7)
Rutgers University students and employees (31)	829	No	No	41 (4.9)	27 (65.9)
Greek citizens evacuated from the United Kingdom, Spain, and Turkey (32)†	783	<b>Yes</b>	No	40 (5.1)	35 (87.5)
Boston, Massachusetts, obstetric patients (33)	757	No	No	20 (2.6)	9 (45.0)
Córdoba, Colombia, residents (34)	686	No	No	35 (5.1)	18 (51.4)
New York City obstetric patients 1 (35)	675	No	No	70 (10.4)	55 (78.6)
Santiago, Chile, obstetric patients (36)	586	No	No	37 (6.3)	16 (43.2)
Japanese citizens evacuated from Wuhan, China (37)	564	<b>Yes</b>	No	11 (2.0)	3 (27.3)
London nursing home residents and staff (38)	518	<b>Yes</b>	No	158 (30.5)	72 (45.6)
Indian citizens evacuated from Iran (39)	474	<b>Yes</b>	No	48 (10.1)	44 (91.7)
Maryland long-term care facility residents (40)	426	<b>Yes</b>	No	177 (41.5)	154 (87.0)
South India retinal surgery patients (41)	413	No	No	9 (2.2)	9 (100.0)
Boston homeless shelter occupants (42)	408	No	No	147 (36.0)	129 (87.8)
Seafood plant employees (43)§	376	No	No	124 (33.0)	118 (95.0)
Genoa, Italy, obstetric patients (44)	333	No	No	7 (2.1)	6 (85.7)
London maternity hospital staff (45)	266	<b>Yes</b>	No	47 (17.7)	16 (34.0)
Argentine cruise ship passengers and crew (46)	217	No	No	128 (59.0)	104 (81.3)
New York City obstetric patients 2 (47)	214	<b>Yes</b>	No	33 (15.4)	29 (87.9)
Bogotá, Colombia, airport employees (48)	212	<b>Yes</b>	No	35 (16.5)	24 (68.6)
Porto, Portugal, obstetric patients (49)	184	No	No	11 (6.0)	9 (81.8)
Los Angeles, California, homeless shelter occupants (50)	178	No	No	43 (24.2)	27 (62.8)
Illinois skilled-nursing facility residents (51)	126	<b>Yes</b>	No	33 (26.2)	13 (39.4)
Boston grocery store employees (52)	104	No	No	21 (20.2)	16 (76.2)
Los Angeles skilled-nursing facility residents (53)	99	<b>Yes</b>	No	19 (19.2)	6 (31.6)
King County nursing facility residents (54)	76	<b>Yes</b>	No	48 (63.2)	3 (6.3)

PCR = polymerase chain reaction; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

\* Boldface indicates details that increase the likelihood of higher-quality evidence.

† Data clarified via personal communication with coauthor.

‡ Percentages reflect weighting by the study's authors to estimate statewide prevalence.

§ Estimated from incomplete source data.

090177e1981d7da1\Final\Final On: 20-Sep-2021 20:00 (GMT)

**Table 2. Antibody Testing**

Study or Report	Tested, n*	Random Sampling*	SARS-CoV-2 Positive, n (%)	Asymptomatic, n (%)
England residents (55)	<b>365 104</b>	<b>Yes</b>	17 576 (4.8)	5694 (32.4)
Spain residents (56)	<b>61 075</b>	<b>Yes</b>	3053 (5.0)	1008 (33.0)
Detroit, Michigan, hospital staff (57)	<b>20 614</b>	No	1818 (8.8)	798 (43.9)
Wuhan, China, hospital staff (58)	8553	No	424 (5.0)	148 (34.9)
Bavaria, Germany, children aged 1-18 y (59)	4859	<b>Yes</b>	47 (1.0)	22 (46.8)
Louisiana residents (60)	4778	<b>Yes</b>	311 (6.5)	147 (47.3)
Munich, Germany, hospital staff (61)	4554	No	108 (2.4)	28 (25.9)
Cairo, Egypt, hospital staff (62)	4040	No	170 (4.2)	116 (68.2)
Health care personnel at 13 U.S. medical centers (63)	3248	No	194 (6.0)	56 (28.9)
Maranhão, Brazil, residents (64)	3156	<b>Yes</b>	1167 (37.0)	320 (27.4)
Ischgl, Austria, residents (65)	1473	No	622 (42.2)	529 (85.0)
Wuhan dialysis patients (66)	1027	No	99 (9.6)	50 (50.5)
Buenos Aires, Argentina, residents (67)	873	No	466 (53.4)	396 (85.0)
Connecticut residents (68)	567	<b>Yes</b>	23 (4.1)	5 (21.7)
Sweden nursing home staff (69)	459	No	86 (18.7)	40 (46.5)
London, England, dialysis patients (70)	356	No	129 (36.2)	52 (40.3)
Nashville, Tennessee, hospital staff (71)	249	No	19 (7.6)	8 (42.1)
London maternity unit staff (72)	200	No	29 (14.5)	10 (34.5)

SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

\* Boldface indicates details that increase the likelihood of higher-quality evidence.

“Among confirmed SARS CoV 2 infections, we did not observe significant differences in the frequency of asymptomatic infection between age groups” (17).

Of the 43 studies that used PCR testing, 21 involved high density living or working environments, such as nursing homes and factories (13, 15, 18, 19, 21, 22, 24, 28, 30, 38, 40, 42, 46, 50, 51, 53, 54). The settings with the highest proportion of persons who tested positive without symptoms included prisons (19) and poultry processing plants (21). Yet, the data seem to be insufficient to conclude that setting was a causative factor. In the 21 studies of high density environments, the proportion of persons who tested positive but had no symptoms at the time of testing ranged from 6.3% to 96.0%, with a median of 62.8% (IQR, 40.6% to 87.0%). In the remaining 22 studies that did not involve such high density environments, the proportion ranged from 27.3% to 100%, with a median of 67.2% (IQR, 43.5% to 84.7%).

**Antibody Testing**

In the 18 studies based on antibody testing (Table 2) (55-72), the proportion of persons who tested positive but did not report having had symptoms ranged from 21.7% to 85.0%, with a median of 41.2% (IQR, 32.6% to 48.1%).

Among the 18 antibody testing studies, 6 used random selection of participants to achieve a representative sample of their target population: residents of England (55); Spain (56); Bavaria, Germany (59); Louisiana (60); Maranhão, Brazil (64); or Connecticut (68). In these antibody studies with representative samples, the proportion of persons who tested positive but did not report having had symptoms ranged from 21.7% to 47.3%, with a median of 32.7% (IQR, 28.7% to 43.4%).

The 2 largest studies based on antibody testing were nationwide serosurveys from England (55) and Spain (56), both designed to achieve representative samples of

**Table 3. Evolution of Symptomatic Status**

Study	Initially Tested Positive Without Symptoms, n	Remained Asymptomatic, n (%)
U.S. skilled-nursing facility residents (15)	3227	2194 (68.0)
Vo', Italy, residents (17)	34	34 (100.0)
U.S. Navy aircraft carrier crew (18)	978	572 (58.5)
San Francisco, California, residents (20)	41	23 (56.1)
Diamond Princess cruise ship passengers and crew (22)	410	311 (75.9)
Greek citizens evacuated from the United Kingdom, Spain, and Turkey (32)*	39	35 (89.7)
Japanese citizens evacuated from Wuhan, China (37)	6	3 (50.0)
London, England, nursing home residents and staff (38)	67	46 (68.7)
Indian citizens evacuated from Iran (39)	44	44 (100.0)
Maryland long-term care facility residents (40)	177	154 (87.0)
New York City obstetric patients 2 (47)	29	26 (89.7)
Illinois skilled-nursing facility residents (51)	14	13 (92.9)
Los Angeles, California, skilled-nursing facility residents (53)	14	6 (42.9)
King County, Washington, nursing facility residents (54)	27	3 (11.1)

\* Data clarified via personal communication with coauthor.

090177e1981d7da1\Final\Final On: 20-Sep-2021 20:00 (GMT)

community dwelling persons. The English data, from the REACT program described earlier, were collected during 3 rounds of testing from June through September 2020 and include 365 104 persons. The Spanish data were collected 27 April to 11 May 2020 and include 61 075 persons. The proportion of persons who tested positive but did not report having had symptoms was 32.4% in England and 33.0% in Spain.

## DISCUSSION

Symptom detection relies on the subjective reports of patients (73). For example, anosmia has turned out to be a distinctive symptom of COVID 19 (74), and we depend on patients to perceive and report a diminution, however slight, of their normal olfactory abilities. But such self reports are influenced by many factors, including variability in the ability to recall symptoms and idiosyncratic awareness of bodily sensations.

Current data suggest that infected persons without symptoms including both presymptomatic and asymptomatic persons account for more than 40% of all SARS CoV 2 transmission (75 77). The proportion of new infections caused by asymptomatic persons alone is uncertain, but when researchers in Wanzhou, China, analyzed epidemiologic data for “183 confirmed COVID 19 cases and their close contacts from five generations of transmission,” they determined that the asymptomatic cases, which made up 32.8% of infected persons, caused 19.3% of infections (78).

The 61 studies and reports that we have collected provide compelling evidence that the asymptomatic fraction of SARS CoV 2 infection is sizable. These data enable us to make reasonable inferences about the proportion of SARS CoV 2 infections that are asymptomatic.

Studies designed to achieve representative samples of large populations provide useful data because they may accurately reflect human populations in general. Four of the PCR based studies are in this category, with target populations of England (10 12, 14), Iceland (16), and Indiana (23). The proportion of persons who tested positive but had no symptoms at the time of testing ranged from 43.0% to 76.5%, with a median of 45.6% (IQR, 43.6% to 61.8%). However, these studies fall short of providing the highest quality evidence because they collected only cross sectional data. As a result, we cannot distinguish between presymptomatic and asymptomatic cases.

In 14 longitudinal studies that reported information on the evolution of symptomatic status, a median of 72.3% of persons who tested positive but had no symptoms at the time of testing remained asymptomatic during a follow up period (15, 17, 18, 20, 22, 32, 37 40, 47, 51, 53, 54). If a similar proportion remained asymptomatic in the 4 large, representative, PCR based studies, in which the median was 45.6%, the asymptomatic fraction of SARS CoV 2 infection would be 33.0%.

Among the data that we have assembled here, the highest quality evidence comes from the large scale studies using antibody testing that were designed to achieve representative samples of nationwide populations in

England ( $n = 365\,104$ ) (55) and Spain ( $n = 61\,075$ ) (56). It is remarkable that these independently conducted serosurveys yielded nearly identical proportions of asymptomatic SARS CoV 2 infections: 32.4% in England and 33.0% in Spain.

We may infer that persons who receive positive antibody test results can be classified accurately as asymptomatic because such results are likely to occur only after the onset of symptoms, if any. In a study of 222 hospitalized patients in Wuhan, China, IgM and IgG antibodies to SARS CoV 2 were first detected 3 and 4 days, respectively, after symptomatic onset of COVID 19 (79). In a study of 109 health care workers and 64 hospitalized patients in Zurich, Switzerland, the severity of illness seemed to affect how quickly SARS CoV 2 antibodies appeared (80). Patients with severe COVID 19 had detectable SARS CoV 2 antibody titers after symptom onset, but those with mild cases “remained negative or became positive [for SARS CoV 2 antibodies] 12 to 14 days after symptom onset” (80). These data suggest that positive antibody test results are unlikely to occur during the period when it is uncertain whether an infected person is presymptomatic or asymptomatic.

However, serosurveys do have significant limitations for the purpose of estimating the asymptomatic fraction. Not all persons who are believed to have been infected with SARS CoV 2 later have a positive result for SARS CoV 2 antibodies (81). The reasons may include a false positive result on the initial PCR test; a false negative result on the antibody test; or the absence of detectable antibodies, perhaps because the infection was cleared without requiring adaptive immunity. In addition, the role of mucosal immunity in clearing SARS CoV 2 infection has not yet been fully elucidated (82), and a nasal wash to detect the IgA antibodies active in mucosal immunity is not part of standard testing practice. Persons who clear SARS CoV 2 infection through innate or mucosal immunity might be more likely to be asymptomatic but would not be categorized as such in a serosurvey, possibly contributing to an underestimate of the asymptomatic fraction.

Another limitation of serosurveys is the requirement that an interview or questionnaire about symptomatic status accompany the blood sample. The onus is on the study participant to accurately recall symptoms, if any, from weeks or even months earlier. In the midst of a pandemic that has transformed everyday life around the globe, it seems reasonable to hypothesize that awareness of and memory for symptoms possibly related to COVID 19 are heightened. This might result in a greater likelihood of noticing and reporting symptoms that would otherwise be missed or ignored, thereby leading to a lower estimate of the asymptomatic fraction. For these reasons, we have evaluated serosurveys in the context of other results and found them to be concordant.

When estimates from large scale, cross sectional, PCR based studies with representative samples; longitudinal PCR based studies; and nationwide serosurveys with representative samples are combined, it seems that the asymptomatic fraction of SARS CoV 2 infection is at least one third. To confirm this estimate, large scale

longitudinal studies using PCR testing with representative samples of national populations would be useful. As SARS CoV 2 vaccination campaigns are implemented worldwide, though, the window for such research may be closing.

In light of the data presented here, we believe that COVID 19 control strategies must be altered, taking into account the prevalence and transmission risk of asymptomatic SARS CoV 2 infection. Frequent, inexpensive, rapid home tests (83) to identify and contain presymptomatic or asymptomatic cases along with government programs that provide financial assistance and, if necessary, housing to enable infected persons to isolate themselves (84) may be a viable option. And as the first generation of SARS CoV 2 vaccines is deployed, more research will be needed to determine their efficacy in preventing asymptomatic infection (85).

From Scripps Research Translational Institute, La Jolla, California (D.P.O., E.J.T.).

**Grant Support:** By grant UL1TR002550 from the National Institutes of Health.

**Disclosures:** Authors have disclosed no conflicts of interest. Forms can be viewed at [www.acponline.org/authors/icmje/ConflictOfInterestForms.do?msNum\\_M20\\_6976](http://www.acponline.org/authors/icmje/ConflictOfInterestForms.do?msNum_M20_6976).

**Reproducible Research Statement:** *Study protocol and statistical code:* Not applicable. *Data set:* All of the data on which the authors based their analysis have been published with this review.

**Corresponding Author:** Eric J. Topol, MD, Scripps Research Translational Institute, 3344 North Torrey Pines Court, 3rd Floor, La Jolla, CA 92037; e mail, [etopol@scripps.edu](mailto:etopol@scripps.edu).

Current author addresses and author contributions are available at [Annals.org](http://Annals.org).

## References

1. Champredon D, Moghadas SM. Quantifying the contribution of asymptomatic infection to the cumulative incidence. *Epidemiol Infect.* 2017;145:1256-1258. [PMID: 28166860] doi:10.1017/S0950268817000115
2. Centers for Disease Control and Prevention. Measles. Accessed at [www.cdc.gov/vaccines/pubs/pinkbook/meas.html](http://www.cdc.gov/vaccines/pubs/pinkbook/meas.html) on 24 November 2020.
3. Centers for Disease Control and Prevention. About cytomegalovirus (CMV). Accessed at [www.cdc.gov/cmvid/overview.html](http://www.cdc.gov/cmvid/overview.html) on 24 November 2020.
4. European Center for Disease Prevention and Control. Disease factsheet about poliomyelitis. Updated 11 July 2018. Accessed at [www.ecdc.europa.eu/en/poliomyelitis/facts](http://www.ecdc.europa.eu/en/poliomyelitis/facts) on 3 October 2020.
5. Oran DP, Topol EJ. Prevalence of asymptomatic SARS CoV 2 infection. A narrative review. *Ann Intern Med.* 2020;173:362-367. doi:10.7326/M20\_3012
6. Apuzzo M, Gebrekidan S, Kirkpatrick DD. How the world missed COVID 19's silent spread. *New York Times.* 27 June 2020. Accessed at [www.nytimes.com/2020/06/27/world/europe/coronavirus-spread-asymptomatic.html](http://www.nytimes.com/2020/06/27/world/europe/coronavirus-spread-asymptomatic.html) on 1 October 2020.

7. Lauer SA, Grantz KH, Bi Q, et al. The incubation period of coronavirus disease 2019 (COVID 19) from publicly reported confirmed cases: estimation and application. *Ann Intern Med.* 2020;172:577-582. doi:10.7326/M20\_0504
8. Centers for Disease Control and Prevention. Interim clinical guidance for management of patients with confirmed coronavirus disease (COVID 19). 10 September 2020. Accessed at [www.cdc.gov/coronavirus/2019-ncov/hcp/clinical-guidance-management-patients.html](http://www.cdc.gov/coronavirus/2019-ncov/hcp/clinical-guidance-management-patients.html) on 4 October 2020.
9. Furukawa NW, Brooks JT, Sobel J. Evidence supporting transmission of severe acute respiratory syndrome coronavirus 2 while presymptomatic or asymptomatic. *Emerg Infect Dis.* 2020;26. [PMID: 32364890] doi:10.3201/eid2607.201595
10. Riley S, Ainslie KEC, Eales O, et al. High prevalence of SARS CoV 2 swab positivity in England during September 2020: interim report of round 5 of REACT 1 study. medRxiv. Preprint posted online 2 October 2020. doi:10.1101/2020.09.30.20204727
11. Riley S, Ainslie KEC, Eales O, et al. High and increasing prevalence of SARS CoV 2 swab positivity in England during end September beginning October 2020: REACT 1 round 5 updated report. medRxiv. Preprint posted online 14 October 2020. doi:10.1101/2020.10.12.20211227
12. Riley S, Ainslie KEC, Eales O, et al. REACT 1 round 6 updated report: high prevalence of SARS CoV 2 swab positivity with reduced rate of growth in England at the start of November 2020. 12 November 2020. Accessed at <https://spiral.imperial.ac.uk/handle/10044/1/83912> on 26 November 2020.
13. Hoxha A, Wyndham Thomas C, Klamer S, et al. Asymptomatic SARS CoV 2 infection in Belgian long term care facilities [Letter]. *Lancet Infect Dis.* 2020. [PMID: 32628906] doi:10.1016/S1473-3099(20)30560-0
14. Petersen I, Phillips A. Three quarters of people with SARS CoV 2 infection are asymptomatic: analysis of English household survey data. *Clin Epidemiol.* 2020;12:1039-1043. [PMID: 33116898] doi:10.2147/CLEP.S276825
15. White EM, Santostefano CM, Feifer RA, et al. Asymptomatic and presymptomatic severe acute respiratory syndrome coronavirus 2 infection rates in a multistate sample of skilled nursing facilities. *JAMA Intern Med.* 2020;180:1709-1711. [PMID: 33074318] doi:10.1001/jamainternmed.2020.5664
16. Gudbjartsson DF, Helgason A, Jonsson H, et al. Spread of SARS CoV 2 in the Icelandic population. *N Engl J Med.* 2020;382:2302-2315. [PMID: 32289214] doi:10.1056/NEJMoa2006100
17. Lavezzo E, Franchin E, Ciavarella C, et al; Imperial College COVID 19 Response Team. Suppression of a SARS CoV 2 outbreak in the Italian municipality of Vo'. *Nature.* 2020;584:425-429. [PMID: 32604404] doi:10.1038/s41586-020-2488-1
18. Kasper MR, Geibe JR, Sears CL, et al. An outbreak of Covid 19 on an aircraft carrier. *N Engl J Med.* 2020;383:2417-2426. [PMID: 33176077] doi:10.1056/NEJMoa2019375
19. So L, Smith G. In four U.S. state prisons, nearly 3,300 inmates test positive for coronavirus 96% without symptoms. *Reuters.* 25 April 2020. Accessed at [www.reuters.com/article/us-health-coronavirus-prisons-testing/idUSKCN2270RX](http://www.reuters.com/article/us-health-coronavirus-prisons-testing/idUSKCN2270RX) on 26 April 2020.
20. Chamie G, Marquez C, Crawford E, et al; CLIAhub Consortium. SARS CoV 2 community transmission disproportionately affects Latinx population during shelter in place in San Francisco. *Clin Infect Dis.* 2020. [PMID: 32821935] doi:10.1093/cid/ciaa1234
21. Lush T. Hundreds test positive at Tyson Foods plant in Arkansas. *Associated Press.* 21 June 2020. Accessed at <https://apnews.com/article/9b2b24d3b3ed981041a7fe6ab499100c> on 6 October 2020.
22. Sakurai A, Sasaki T, Kato S, et al. Natural history of asymptomatic SARS CoV 2 infection [Letter]. *N Engl J Med.* 2020;383:885-886. [PMID: 32530584] doi:10.1056/NEJMc2013020
23. Menachemi N, Yiannoutsos CT, Dixon BE, et al. Population point prevalence of SARS CoV 2 infection based on a statewide

- random sample Indiana, April 25-29, 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69:960-964. [PMID: 32701938] doi:10.15585/mmwr.mm6929e1
24. Marossy A, Rakowicz S, Bhan A, et al. A study of universal SARS CoV 2 RNA testing of residents and staff in a large group of care homes in South London. *J Infect Dis.* 2020. [PMID: 32889532] doi:10.1093/infdis/jiaa565
  25. Letizia AG, Ramos I, Obla A, et al. SARS CoV 2 transmission among Marine recruits during quarantine. *N Engl J Med.* 2020; 383:2407-2416. [PMID: 33176093] doi:10.1056/NEJMoa2029717
  26. Centre d'Épidémiologie et de Santé Publique des Armées. Investigation de l'épidémie de COVID 19 au sein du Groupe Aéronaval: 21 janvier - 13 avril 2020. 4 May 2020. Accessed at [www.defense.gouv.fr/content/download/583466/9938746/file/20200405\\_929\\_ARM\\_SSA\\_CESPA\\_rapport\\_epidemie\\_covid19\\_Gan\\_VEXP.pdf](http://www.defense.gouv.fr/content/download/583466/9938746/file/20200405_929_ARM_SSA_CESPA_rapport_epidemie_covid19_Gan_VEXP.pdf) on 4 June 2020.
  27. Ly TDA, Zanini D, Laforge V, et al. Pattern of SARS CoV 2 infection among dependant elderly residents living in long term care facilities in Marseille, France, March-June 2020. *Int J Antimicrob Agents.* 2020;56:106219. [PMID: 33189890] doi:10.1016/j.ijantimicag.2020.106219
  28. Rogers JH, Link AC, McCulloch D, et al. Characteristics of COVID 19 in homeless shelters: a community based surveillance study. *Ann Intern Med.* 2020. doi:10.7326/M20-3799
  29. Hempel D, Milani V, Kleespies A, et al. SARS CoV 2 infections in outpatients with cancer: most infected patients are asymptomatic carriers without impact on chemotherapy [Abstract]. *Ann Oncol.* 2020;31(Suppl 4):S995. Abstract no. 1680P. doi:10.1016/j.annonc.2020.08.1744
  30. Feaster M, Goh YY. High proportion of asymptomatic SARS CoV 2 infections in 9 long term care facilities, Pasadena, California, USA, April 2020. *Emerg Infect Dis.* 2020;26:2416-2419. [PMID: 32614768] doi:10.3201/eid2610.202694
  31. Barrett ES, Horton DB, Roy J, et al. Prevalence of SARS CoV 2 infection in previously undiagnosed health care workers at the onset of the U.S. COVID 19 epidemic. *medRxiv.* 2020. [PMID: 32511600] doi:10.1101/2020.04.20.20072470
  32. Lytras T, Dellis G, Flountzi A, et al. High prevalence of SARS CoV 2 infection in repatriation flights to Greece from three European countries. *J Travel Med.* 2020;27. [PMID: 32297940] doi:10.1093/jtm/taaa054
  33. Goldfarb IT, Diouf K, Barth WH, et al. Universal SARS CoV 2 testing on admission to the labor and delivery unit: low prevalence among asymptomatic obstetric patients. *Infect Control Hosp Epidemiol.* 2020;41:1095-1096. [PMID: 32456729] doi:10.1017/ice.2020.255
  34. Mattar S, Martinez Bravo C, Rivero R, et al. Epidemiological and viral features of a cohort of SARS CoV 2 symptomatic and asymptomatic individuals in an area of the Colombian Caribbean. *Research Square.* Preprint posted online 30 October 2020. doi:10.21203/rs.3.rs-57254/v3
  35. Prabhu M, Cagino K, Matthews KC, et al. Pregnancy and post partum outcomes in a universally tested population for SARS CoV 2 in New York City: a prospective cohort study. *BJOG.* 2020; 127:1548-1556. [PMID: 32633022] doi:10.1111/1471-0528.16403
  36. Díaz Corvillón P, Mönckeberg M, Barros A, et al. Routine screening for SARS CoV 2 in unselected pregnant women at delivery. *PLoS One.* 2020;15:e0239887. [PMID: 32991621] doi:10.1371/journal.pone.0239887
  37. Kutsuna S, Suzuki T, Hayakawa K, et al. SARS CoV 2 screening test for Japanese returnees from Wuhan, China, January 2020. *Open Forum Infect Dis.* 2020;7:ofaa243. [PMID: 32754627] doi:10.1093/ofid/ofaa243
  38. Ladhani SN, Chow JY, Janarthanan R, et al. Investigation of SARS CoV 2 outbreaks in six care homes in London, April 2020. *EclinicalMedicine.* 2020;26:100533. [PMID: 32923993] doi:10.1016/j.eclinm.2020.100533
  39. Saurabh S, Kumar R, Kumar N, et al. Dynamics of SARS CoV 2 transmission among Indian nationals evacuated from Iran. *Disaster Med Public Health Prep.* 2020;1:7. [PMID: 33087205] doi:10.1017/dmp.2020.393
  40. Bigelow BF, Tang O, Barshick B, et al. Outcomes of universal COVID 19 testing following detection of incident cases in 11 long term care facilities. *JAMA Intern Med.* 2021;181:127-129. [PMID: 32662818] doi:10.1001/jamainternmed.2020.3738
  41. Kannan NB, Sen S, Reddy H, et al. Preoperative COVID 19 testing for elective vitreoretinal surgeries: experience from a major tertiary care institute in South India. *Indian J Ophthalmol.* 2020;68: 2373-2377. [PMID: 33120621] doi:10.4103/ijo.IJO-2870-20
  42. Baggett TP, Keyes H, Sporn N, et al. COVID 19 outbreak at a large homeless shelter in Boston: implications for universal testing. *medRxiv.* Preprint posted online 15 April 2020. doi:10.1101/2020.04.12.20059618
  43. Cline S. Cases at seafood plant cause spike in Oregon COVID numbers. *Associated Press.* 9 June 2020. Accessed at <https://apnews.com/article/4b9d38f206db9ce5267a5898ac24f238> on 6 October 2020.
  44. Massarotti C, Adriano M, Cagnacci A, et al. Asymptomatic SARS CoV 2 infections in pregnant patients in an Italian city during the complete lockdown. *J Med Virol.* 2020. [PMID: 32841411] doi:10.1002/jmv.26458
  45. Khalil A, Hill R, Ladhani S, et al. COVID 19 screening of health care workers in a London maternity hospital [Letter]. *Lancet Infect Dis.* 2021;21:23-24. [PMID: 32437699] doi:10.1016/S1473-3099(20)30403-5
  46. Ing AJ, Cocks C, Green JP. COVID 19: in the footsteps of Ernest Shackleton. *Thorax.* 2020;75:693-694. [PMID: 32461231] doi:10.1136/thoraxjnl-2020-215091
  47. Sutton D, Fuchs K, D'Alton M, et al. Universal screening for SARS CoV 2 in women admitted for delivery [Letter]. *N Engl J Med.* 2020;382:2163-2164. [PMID: 32283004] doi:10.1056/NEJMc2009316
  48. Malagón Rojas J, Gómez Rendón C, Parra EL, et al. SARS CoV 2 and RT-PCR in asymptomatic patients: results of a cohort of workers at El Dorado International Airport in Bogotá, 2020. *Biomedica.* 2020;40:166-172. [PMID: 33152200] doi:10.7705/biomedica.5802
  49. Figueiredo R, Tavares S, Moucho M, et al. Systematic screening for SARS CoV 2 in pregnant women admitted for delivery in a Portuguese maternity. *J Perinat Med.* 2020;48:977-980. [PMID: 33085639] doi:10.1515/jpm-2020-0387
  50. Chou E. Dozens positive for coronavirus at LA's Skid Row homeless shelter, after all residents tested. *Los Angeles Daily News.* 21 April 2020. Updated 22 April 2020. Accessed at [www.dailynews.com/2020/04/21/dozens-positive-for-coronavirus-at-las-skid-row-homeless-shelter-after-all-residents-tested-on-23-april-2020](http://www.dailynews.com/2020/04/21/dozens-positive-for-coronavirus-at-las-skid-row-homeless-shelter-after-all-residents-tested-on-23-april-2020).
  51. Patel MC, Chaisson LH, Borgetti S, et al. Asymptomatic SARS CoV 2 infection and COVID 19 mortality during an outbreak investigation in a skilled nursing facility. *Clin Infect Dis.* 2020;71:2920-2926. [PMID: 32548628] doi:10.1093/cid/ciaa763
  52. Lan FY, Suharlilim C, Kales SN, et al. Association between SARS CoV 2 infection, exposure risk and mental health among a cohort of essential retail workers in the USA. *Occup Environ Med.* 2020. [PMID: 33127659] doi:10.1136/oemed-2020-106774
  53. Dora AV, Winnett A, Jatt LP, et al. Universal and serial laboratory testing for SARS CoV 2 at a long term care skilled nursing facility for veterans - Los Angeles, California, 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69:651-655. [PMID: 32463809] doi:10.15585/mmwr.mm6921e1
  54. Arons MM, Hatfield KM, Reddy SC, et al; Public Health Seattle and King County and CDC COVID 19 Investigation Team. Presymptomatic SARS CoV 2 infections and transmission in a skilled nursing facility. *N Engl J Med.* 2020;382:2081-2090. [PMID: 32329971] doi:10.1056/NEJMoa2008457

55. Ward H, Cooke G, Atchison C, et al. Declining prevalence of antibody positivity to SARS CoV 2: a community study of 365,000 adults. medRxiv. Preprint posted online 27 October 2020. doi:10.1101/2020.10.26.20219725
56. Pollán M, Pérez Gómez B, Pastor Barriuso R, et al; ENE COVID Study Group. Prevalence of SARS CoV 2 in Spain (ENE COVID): a nationwide, population based seroepidemiological study. *Lancet*. 2020;396:535-544. [PMID: 32645347] doi:10.1016/S0140-6736(20)31483-5
57. Sims MD, Maine GN, Childers KL, et al; BLAST COVID 19 Study Group. COVID 19 seropositivity and asymptomatic rates in health care workers are associated with job function and masking. *Clin Infect Dis*. 2020. [PMID: 33150375] doi:10.1093/cid/ciaa1684
58. Zhang S, Guo M, Wu F, et al. Factors associated with asymptomatic infection in health care workers with severe acute respiratory syndrome coronavirus 2 infection in Wuhan, China: a multi-centre retrospective cohort study. *Clin Microbiol Infect*. 2020;26:1670-1675. [PMID: 32911080] doi:10.1016/j.cmi.2020.08.038
59. Hippich M, Holthaus L, Assfalg R, et al. Public health antibody screening indicates a six fold higher SARS CoV 2 exposure rate than reported cases in children. *Med (N Y)*. 2020. [PMID: 33163984] doi:10.1016/j.medj.2020.10.003
60. Feehan A, Fort D, Garcia Diaz J, et al. Frequency of symptoms and asymptomatic SARS CoV 2 infection in Louisiana, USA. SSRN. Preprint posted online 2 October 2020. doi:10.2139/ssrn.3702933
61. Erber J, Kappler V, Haller B, et al. Strategies for infection control and prevalence of anti SARS CoV 2 IgG antibodies in 4,554 employees of a university hospital in Munich, Germany. SSRN. Preprint posted online 13 October 2020. doi:10.2139/ssrn.3678599
62. Mostafa A, Kandil S, El Sayed MH, et al. Universal COVID 19 screening of 4040 health care workers in a resource limited setting: an Egyptian pilot model in a university with 12 public hospitals and medical centers. *Int J Epidemiol*. 2020. [PMID: 33094320] doi:10.1093/ije/dyaa173
63. Self WH, Tenforde MW, Stubblefield WB, et al; CDC COVID 19 Response Team. Seroprevalence of SARS CoV 2 among frontline health care personnel in a multistate hospital network – 13 academic medical centers, April–June 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69:1221-1226. [PMID: 32881855] doi:10.15585/mmwr.mm6935e2
64. Silva AMA, Lima Neto LG, Azevedo CMPS, et al. Population based seroprevalence of SARS CoV 2 is more than halfway through the herd immunity threshold in the State of Maranhao, Brazil. medRxiv. Preprint posted online 1 September 2020. doi:10.1101/2020.08.28.20180463
65. Österreichischer Rundfunk. Ischgl: 42,4 Prozent haben Antikörper. 25 June 2020. Accessed at <https://tirol.orf.at/stories/3054826> on 5 October 2020.
66. Tang H, Tian JB, Dong JW, et al. Serologic detection of SARS CoV 2 infections in hemodialysis centers: a multicenter retrospective study in Wuhan, China. *Am J Kidney Dis*. 2020;76:490-499.e1. [PMID: 32628990] doi:10.1053/j.ajkd.2020.06.008
67. Figar S, Pagotto V, Luna L, et al. Community level SARS CoV 2 seroprevalence survey in urban slum dwellers of Buenos Aires City, Argentina: a participatory research. medRxiv. Preprint posted online 18 July 2020. doi:10.1101/2020.07.14.20153858
68. Mahajan S, Srinivasan R, Redlich CA, et al. Seroprevalence of SARS CoV 2 specific IgG antibodies among adults living in Connecticut: post infection prevalence (PIP) study. *Am J Med*. 2020. [PMID: 33130124] doi:10.1016/j.amjmed.2020.09.024
69. Lindahl JF, Hoffman T, Esmaeilzadeh M, et al. High seroprevalence of SARS CoV 2 in elderly care employees in Sweden. *Infect Ecol Epidemiol*. 2020;10:1789036. [PMID: 32939231] doi:10.1080/2008686.2020.1789036
70. Clarke C, Prendecki M, Dhutia A, et al. High prevalence of asymptomatic COVID 19 infection in hemodialysis patients detected using serologic screening. *J Am Soc Nephrol*. 2020;31:1969-1975. [PMID: 32732391] doi:10.1681/ASN.2020060827
71. Stubblefield WB, Talbot HK, Feldstein L, et al; Influenza Vaccine Effectiveness in the Critically Ill (IVY) Investigators. Seroprevalence of SARS CoV 2 among frontline healthcare personnel during the first month of caring for COVID 19 patients Nashville, Tennessee. *Clin Infect Dis*. 2020. [PMID: 32628750] doi:10.1093/cid/ciaa936
72. Bampoe S, Lucas DN, Neall G, et al. A cross sectional study of immune seroconversion to SARS CoV 2 in frontline maternity health professionals. *Anaesthesia*. 2020;75:1614-1619. [PMID: 32777861] doi:10.1111/anae.15229
73. Knapp TR. Symptom: cause, effect, both, or neither? [Editorial]. *Clin Nurs Res*. 2018;27:391-394. [PMID: 29595078] doi:10.1177/1054773818761904
74. Aziz M, Goyal H, Haghbin H, et al. The association of “loss of smell” to COVID 19: a systematic review and meta analysis. *Am J Med Sci*. 2020. [PMID: 33349441] doi:10.1016/j.amjms.2020.09.017
75. He X, Lau EHY, Wu P, et al. Temporal dynamics in viral shedding and transmissibility of COVID 19. *Nat Med*. 2020;26:672-675. [PMID: 32296168] doi:10.1038/s41591-020-0869-5
76. Ma S, Zhang J, Zeng M, et al. Epidemiological parameters of COVID 19: case series study. *J Med Internet Res*. 2020;22:e19994. [PMID: 33001833] doi:10.2196/19994
77. Zhang H, Hong C, Zheng Q, et al. A multi family cluster of COVID 19 associated with asymptomatic and pre symptomatic transmission in Jixi City, Heilongjiang, China, 2020 [Letter]. *Emerg Microbes Infect*. 2020;9:2509-2514. [PMID: 33238813] doi:10.1080/22221751.2020.1837015
78. Shi Q, Hu Y, Peng B, et al. Effective control of SARS CoV 2 transmission in Wanzhou, China. *Nat Med*. 2020. [PMID: 33257893] doi:10.1038/s41591-020-01178-5
79. Zhang B, Zhou X, Zhu C, et al. Immune phenotyping based on the neutrophil to lymphocyte ratio and IgG level predicts disease severity and outcome for patients with COVID 19. *Front Mol Biosci*. 2020;7:157. [PMID: 32719810] doi:10.3389/fmolb.2020.00157
80. Cervia C, Nilsson J, Zurbuchen Y, et al. Systemic and mucosal antibody responses specific to SARS CoV 2 during mild versus severe COVID 19. *J Allergy Clin Immunol*. 2020. [PMID: 33221383] doi:10.1016/j.jaci.2020.10.040
81. Jiang C, Wang Y, Hu M, et al. Antibody seroconversion in asymptomatic and symptomatic patients infected with severe acute respiratory syndrome coronavirus 2 (SARS CoV 2). *Clin Transl Immunology*. 2020;9:e1182. [PMID: 33005417] doi:10.1002/cti2.1182
82. Russell MW, Moldoveanu Z, Ogra PL, et al. Mucosal immunity in COVID 19: a neglected but critical aspect of SARS CoV 2 infection. *Front Immunol*. 2020;11:611337. [PMID: 33329607] doi:10.3389/fimmu.2020.611337
83. Mina MJ, Parker R, Larremore DB. Rethinking Covid 19 test sensitivity – a strategy for containment. *N Engl J Med*. 2020;383:e120. [PMID: 32997903] doi:10.1056/NEJMp2025631
84. Stieg C. Could you get paid to quarantine during the Covid 19 pandemic? Some local governments are already doing it. CNBC. 2 September 2020. Accessed at [www.cnbc.com/2020/09/02/getting-paid-to-quarantine-during-the-covid-19-pandemic.html](http://www.cnbc.com/2020/09/02/getting-paid-to-quarantine-during-the-covid-19-pandemic.html) on 1 December 2020.
85. Bleier BS, Ramanathan M Jr, Lane AP. COVID 19 vaccines may not prevent nasal SARS CoV 2 infection and asymptomatic transmission. *Otolaryngol Head Neck Surg*. 2020;194:599820982633. [PMID: 33320052] doi:10.1177/0194599820982633



**Current Author Addresses:** Mr. Oran and Dr. Topol: Scripps Research Translational Institute, 3344 North Torrey Pines Court, 3rd Floor, La Jolla, CA 92037.

**Author Contributions:** Conception and design: D.P. Oran, E.J. Topol.

Analysis and interpretation of the data: D.P. Oran, E.J. Topol.

Drafting of the article: D.P. Oran, E.J. Topol.

Critical revision of the article for important intellectual content: D.P. Oran, E.J. Topol.

Final approval of the article: D.P. Oran, E.J. Topol.

Statistical expertise: D.P. Oran.

Obtaining of funding: E.J. Topol.

Administrative, technical, or logistic support: D.P. Oran, E.J. Topol.

Collection and assembly of data: D.P. Oran, E.J. Topol.

090177e1981d7da1\Final\Final On: 20-Sep-2021 20:00 (GMT)